



# 19-0-6

COMMERCIAL

## Turf King Fertilizer With Escalade® Weed & Feed

**THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.**

CONTAINS THE ESTERS OF 2,4-D AND FLUROXYPYR WITH DICAMBA ACID  
Weeds contacted by this product gradually die and disappear from the lawn.  
Treats (or covers) 12,500 Square Feet (100' x 125')

**ACTIVE INGREDIENT:**

Isooctyl (2-ethylhexyl) Ester of 2,4-Dichlorophenoxyacetic Acid*.....	0.8650%
1-Methylheptyl Ester of Fluroxypyr:(4-amino-3-5-dichloro-6-fluoro-2-pyridinyl)oxy)acetic Acid, 1-methylheptyl Ester **.....	0.2060%
Dicamba (3,6-Dichloro-o-Anisic Acid)***.....	0.0717%
<b>OTHER INGREDIENTS:</b> .....	<b>98.8573%</b>

**TOTAL:**..... 100.0000%

Isomer Specific Method, Equivalent to:	
*2,4-Dichlorophenoxyacetic Acid.....	0.5740%
**((4-amino-3-5-dichloro-6-fluoro-2-pyridinyl)oxy)acetic Acid.....	0.1430%
***3,6-Dichloro-o-Anisic Acid.....	0.0717%

Fertilizer guaranteed by Andre and Son, Inc.

**DO NOT SELL, DISTRIBUTE OR USE THE PRODUCT IN NASSAU AND SUFFOLK COUNTY IN NEW YORK.**

### KEEP OUT OF REACH OF CHILDREN CAUTION

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300  
For Medical Emergencies Only, Call (877) 325-1840

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing.

### FIRST AID

**IF IN EYES:**

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

**IF ON SKIN OR CLOTHING:**

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 to 20 minutes.
- Call a poison control center or doctor for treatment advice.

### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

All loaders, applicators, and other handlers must wear:

- long-sleeved shirt and long pants,
- shoes plus socks,
- chemical-resistant gloves made of any waterproof material, selection category A.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### USER SAFETY RECOMMENDATIONS

**Users Should:**

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Distributed and Guaranteed by Andre & Son  
17150 State Route 706 • Montrose, PA 18801 • 570-278-1131

### GUARANTEED ANALYSIS

Total Nitrogen (N) ..... 19.00%  
11.40% Urea Nitrogen  
7.60% CSRUN

Potash (K<sub>2</sub>O) ..... 6.00%

Sulfur (S) ..... 0.55%

**Nutrients derived from:**

Sulfur Coated Urea, Urea, Muriate of Potash..

If this product is being sold in a state requiring the publication of information regarding its metal content, information regarding the contents and levels of metals in this product is available at the following website: <http://www.aapfco.org/metals.htm>

### ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

### DIRECTION FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not enter or allow people (or pets) to enter the treated area until dusts have settled. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

This product is a stable mixture of three herbicides. It is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees and Roughs) and similar turf areas.

This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

<i>Alder</i>	<i>Blue vervain</i>	<i>Canada thistle</i>	<i>Creeping jenny</i>
<i>Annual yellow sweet clover</i>	<i>Box elder</i>	<i>Carolina geranium</i>	<i>Crimson clover</i>
<i>Artichoke</i>	<i>Bracted plantain</i>	<i>Carpetweed</i>	<i>Croton</i>
<i>Aster</i>	<i>Brassbuttons</i>	<i>Catchweed bedstraw</i>	<i>Cudweed</i>
<i>Austrian fieldcress</i>	<i>Bristly oxtongue</i>	<i>Catsear</i>	<i>Curly dock</i>
<i>Bedstraw</i>	<i>Broadleaf dock</i>	<i>Catnip</i>	<i>Curly indigo</i>
<i>Beggarick</i>	<i>Broadleaf plantain</i>	<i>Chickweed</i>	<i>Dandelion</i>
<i>Biden</i>	<i>Broomweed</i>	<i>Chicory</i>	<i>Dead nettle</i>
<i>Bindweed</i>	<i>Buckhorn</i>	<i>Cinquefoil</i>	<i>Dock</i>
<i>Bird vetch</i>	<i>Buckhorn plantain</i>	<i>Clover</i>	<i>Dogbane</i>
<i>Bitterweed</i>	<i>Bulbous buttercup</i>	<i>Cockle</i>	<i>Dogfennel</i>
<i>Bitter wintercress</i>	<i>Bull nettle</i>	<i>Cocklebur</i>	<i>Elderberry</i>
<i>Black-eyed Susan</i>	<i>Bull thistle</i>	<i>Coffeebean</i>	<i>English daisy</i>
<i>Black medic</i>	<i>Burdock</i>	<i>Coffeeweed</i>	<i>Fall dandelion</i>
<i>Black mustard</i>	<i>Burning nettle</i>	<i>Common chickweed</i>	<i>False dandelion</i>
<i>Black-seed plantain</i>	<i>Bur ragweed</i>	<i>Common mullein</i>	<i>False flax</i>
<i>Blessed thistle</i>	<i>Burweed</i>	<i>Common sowthistle</i>	<i>False sunflower</i>
<i>Blue lettuce</i>	<i>Buttercup</i>	<i>Corn Chamomile</i>	<i>Fidleneck</i>

<i>Field bindweed</i>	<i>Mexicanweed</i>	<i>Rough cinquefoil</i>	<i>Trailing Crownvetch</i>
<i>Field pansy</i>	<i>Milk vetch</i>	<i>Tumble mustard</i>	<i>Tumble pigweed</i>
<i>Flea bane (daisy)</i>	<i>Milkweed bloodflower</i>	<i>Roundleaved marigold</i>	<i>Tumbleweed</i>
<i>Flixweed</i>	<i>Mugwort</i>	<i>Rush</i>	<i>Velvetleaf</i>
<i>Florida betony</i>	<i>Morningglory</i>	<i>Russian pigweed</i>	<i>Vernice mallow</i>
<i>Florida pusley</i>	<i>Mouseear chickweed</i>	<i>Russian thistle</i>	<i>Veronica</i>
<i>Frenchweed</i>	<i>Musk thistle</i>	<i>St. Johnswort</i>	<i>Vervain</i>
<i>Galinsoga</i>	<i>Mustard</i>	<i>Scarlet pimpernel</i>	<i>Vetch</i>
<i>Garlic mustard</i>	<i>Narrowleaf plantain</i>	<i>Scotch thistle</i>	<i>Virginia buttonweed</i>
<i>Garlic mustard</i>	<i>Narrowleaf vetch</i>	<i>Sheep sorrel</i>	<i>Virginia creeper</i>
<i>Goathead</i>	<i>Nettle</i>	<i>Shepherdsburse</i>	<i>Virginia pepperweed</i>
<i>Goatsbeard</i>	<i>Orange hawkweed</i>	<i>Slender plantain</i>	<i>Wavyleaf bullthistle</i>
<i>Goldenrod</i>	<i>Oxalis</i>	<i>Smallflower galinsoga</i>	<i>Western clematis</i>
<i>Ground ivy</i>	<i>Oxeye daisy</i>	<i>Smartweed</i>	<i>Western salsify</i>
<i>Gunweed</i>	<i>Parsley-piert</i>	<i>Smooth dock</i>	<i>White clover</i>
<i>Hairy bittercress</i>	<i>Parsnip</i>	<i>Smooth pigweed</i>	<i>White mustard</i>
<i>Hairy fleabane</i>	<i>Pearlwort</i>	<i>Sneezeweed</i>	<i>Wild mustard</i>
<i>Hawkweed</i>	<i>Pennycress</i>	<i>Southern wild rose</i>	<i>Wild aster</i>
<i>Healall</i>	<i>Pennywort</i>	<i>Sowthistle</i>	<i>Wild buckwheat</i>
<i>Heartleaf drymary</i>	<i>Peppergrass</i>	<i>Spanishneedle</i>	<i>Wild carrot</i>
<i>Hedge bindweed</i>	<i>Pepperweed</i>	<i>Spatterdock</i>	<i>Wild four-o'clock</i>
<i>Hedge mustard</i>	<i>Pigweed</i>	<i>Speedwell</i>	<i>Wild garlic</i>
<i>Hemp</i>	<i>Pineywoods bedstraw</i>	<i>Spiny Amaranth</i>	<i>Wild geranium</i>
<i>Henbit</i>	<i>Plains coreopsis</i>	<i>Spiny cocklebur</i>	<i>Wild lettuce</i>
<i>Hoary cress</i>	<i>Plantain</i>	<i>Spotted catsear</i>	<i>Wild marigold</i>
<i>Hoary plantain</i>	<i>Poison hemlock</i>	<i>Spotted knapweed</i>	<i>Wild onion</i>
<i>Hoary vervain</i>	<i>Poison ivy</i>	<i>Spotted spurge</i>	<i>Wild parsnip</i>
<i>Honeysuckle</i>	<i>Poison oak</i>	<i>Spurge</i>	<i>Wild radish</i>
<i>Hop clover</i>	<i>Pokeweed</i>	<i>Spurweed</i>	<i>Wild rape</i>
<i>Horsetail</i>	<i>Poorjoe</i>	<i>Stinging nettle</i>	<i>Wild strawberry</i>
<i>Horsetail</i>	<i>Povertyweed</i>	<i>Stinkweed</i>	<i>Wild sweet potato</i>
<i>Indiana mallow</i>	<i>Prickly lettuce</i>	<i>Stitchwort</i>	<i>Wild vetch</i>
<i>Ironweed</i>	<i>Prickly sida</i>	<i>Strawberry clover</i>	<i>Willow</i>
<i>Jewelweed</i>	<i>Primrose</i>	<i>Sumac</i>	<i>Witchweed</i>
<i>Jimsonweed</i>	<i>Prostrate knotweed</i>	<i>Sunflower</i>	<i>Woolly morningglory</i>
<i>Kochia</i>	<i>Prostrate pigweed</i>	<i>Sweet clover</i>	<i>Woodsorrel</i>
<i>Knawel</i>	<i>Prostrate spurge</i>	<i>Tall nettle</i>	<i>Woolly croton</i>
<i>Knotted</i>	<i>Prostrate vervain</i>	<i>Tall vervain</i>	<i>Woolly plantain</i>
<i>Lambsquarter</i>	<i>Puncture vine</i>	<i>Tansy mustard</i>	<i>Wormseed</i>
<i>Lespedeza</i>	<i>Purslane</i>	<i>Tansy ragwort</i>	<i>Yellow</i>
<i>Locoweed</i>	<i>Ragweed</i>	<i>Tanweed</i>	<i>Yellow rocket</i>
<i>Lupine</i>	<i>Red clover</i>	<i>Tarweed</i>	<i>Yellowflower pepperweed</i>
<i>Mallow</i>	<i>Redroot pigweed</i>	<i>Thistle</i>	<i>and other broadleaf weeds.</i>
<i>Marshelder</i>	<i>Red sorrel</i>	<i>Tick trefoil</i>	
<i>Matchweed</i>	<i>Redstem filaree</i>	<i>Toadflax</i>	

### WHEN TO APPLY

Best results are obtained when weeds are growing actively in spring or early fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including water of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed specified rates because damage to turf may occur.

**NOTE:** When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustinegrasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

### HOW TO APPLY

This 50 pound bag should be used on 12,500 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

For Drop Spreaders: First apply two strips at ends. Then apply product along the length of the lawn, overlapping very slightly and shutting off spreader when turning.

For Rotary Spreaders: Apply in a circular path, starting on the outside and working toward the center, overlapping very slightly.

The following settings are approximate for applying contents of this package at a suggested rate of 4 pounds per 1,000 square feet. These settings are furnished only as a guide. Age, spreader condition, operating speed, and unevenness of terrain can cause variation in applied rates. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound per 250 square feet.

## NET WT 50 lb (22.7 kg)

EPA Reg. No. 228-427-47956 • EPA Est. No. 47956-PA-001

## DIRECTION FOR USE (continued)

### SPREADER SETTINGS ONE PASS APPLICATION\*:

<u>MAKE</u>	<u>SETTING</u>
Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	5
Sears Broadcast	3
Sears	5
Scotts Calibrated	4

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

### Limitations

The maximum rate per application is 6 pounds per 1,000 square feet, limited to 2 applications per year. The maximum seasonal rate is 12 pounds per 1,000 square feet, excluding spot treatments.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

### ENVIRONMENTAL BENEFITS

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment.

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.

- 50' x 50' lawn produces enough oxygen for a family of 4 every day
- Reduces Glare & Reduces Noise
- Traps Dust, Smoke Particles & Other Pollutants
- Purifies Water & Modifies Temperature
- Improves Soil Condition & Controls Erosion

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Always use original container to store pesticides in a secured warehouse or storage building. Do not stack more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

**PESTICIDE DISPOSAL:** If container is damaged or if pesticide has leaked, clean up all spilled material. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, or offered to others for their use, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**For nonrefillable bags of granular or dry formulation**

**CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment, then offer for recycling if available, or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

## LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded. (RV020510)

Escalade is a Registered Trademark of Nufarm Americas Inc. All other trademarks are the property of their respective owners.